

돼지 관상동맥 스텐트 재협착 모델에서 Core[®] Stent와 Palmaz-Schatz[®] Stent의 비교

최동훈 · 최승혁 · 조덕규 · 경희두 · 조정래
김중선 · 오성진 · 권승현 · 장양수 · 조승연

Comparison of the Core[®] Stent and Palmaz-Schatz[®] Stent in a Porcine Stent Restenosis Model

Donghoon Choi, MD, Seung-Hyuk Choi, MD, Duk Kyu Cho, MD, Hee Doo Kyung, MD,
Jung Rae Cho, MD, Jung Sun Kim, MD, Sung Jin Oh, MD,
Seung Hyun Kwon, MD, Yangsoo Jang, MD and Seung Yun Cho, MD

Cardiology Division, Yonsei Cardiovascular Hospital and Cardiovascular Research Institute,
College of Medicine, Yonsei University, Seoul, Korea

ABSTRACT

Background and Objectives : The purpose of this study is to compare the new Core[®] stent and Palmaz-Schatz[®] (PS) stent in a porcine coronary stent restenosis model. **Methods :** Twelve pigs underwent balloon injury followed by implantation of oversized, tubular-type Core[®] and PS[®] stents (stent/artery ratio 1.2 : 1) in twenty-four coronary arteries. Quantitative analyses of the initial and follow-up coronary angiograms at 4 weeks after stenting was performed. The extents of injury and the neointimal area were compared between the two stented groups according to morphometric analysis. The stent flexibility and longitudinal straightening effect were compared between the two groups by the bending test and measurement of the angle changes. **Results :** 1) The reference vessel diameter, stented artery diameter, and diameter of the stenosis were not different between the two groups. 2) The neointimal area was significantly smaller in the Core[®] stent group than in the PS[®] stent group ($1.81 \pm 0.67 \text{ mm}^2$ vs $2.93 \pm 0.94 \text{ mm}^2$, $p = 0.006$). 3) The Core[®] stent had more flexible property than the PS[®] stent. 4) The angle changes following stent implantation did not differ between the two groups (13.2 ± 9.0 , 14.4 ± 11.1 , $p = 0.88$). **Conclusion :** Core[®] stent is effective in the inhibition of neointimal formation in a porcine coronary stent restenosis model. These results may be due to the improved flexibility of the Core[®] stent, although further clinical trials may be needed. (**Korean Circulation J 2001; 31(7):655-661**)

KEY WORDS : Porcine model · Stents · Restenosis.

: 2001 2 28
: 2001 5 28
: , 120 - 752 134
: (02) 361 - 7071 · : (02) 393 - 2041 E - mail : ysjang1212@yumc.yonsei.ac.kr




Fig. 1. Core® stent design permits independent tailoring of radial strength and flexibility elements for optimal stent performance. The thickness of vertical strut is 0.125 mm and the thickness of horizontal strut is 0.09 mm. The cell width of distal portion of stent is 1.2 mm but the width of proximal portion of stent is 1.8 mm.

서론

stent, suboptimal result¹⁻³⁾,
4)5) stent가, tube coil case
Stent, stent, tortuous, angulated, distal le-
sion, slotted tube, stent, coil, stent
tube, radial force가, MAC® stent, multicell-
ular design, slotted tube, stent (flexibility)
vessel support가, 6) MAC®
stent, multicellular design, Core®
stent(E.O. Medical), slotted tube
stent, Palmaz-Schatz®(PS) stent(Johnson & Jo-
hnson), (Fig. 1).

대상 및 방법

실험동물

12, 25, 30 kg, 3, 500 mg, 500 mg, 12, ketamine, xyla-




Fig. 2. The method of measurement of the angle changes. On the orthogonal view the acute angle was measured by a protractor.

zine

8 F sheath, 10,000 IU, MCA - 501® fluoroscope(Medison - ACOMA Co., Seoul, Ko-
rea), 8 F, 스텐트 삽입술 및 추적 관상동맥 조영술, 3.0 mm

1.2 : 1

6, 8, 20

가, Core® stent(1), PS®, stent(2), 1, 12, sheath

Longitudinal straightening effect 측정을 위한 bending test

Core® stent (flexibility) longitudinal straightening effect Palmaz - Schatz® stent

stent PS® stent가 7 mm 1 mm 7 mm PS® stent Core® stent 5 mm

Table 1. Quantitative coronary angiographic findings of porcine coronary arteries after placement of Core® (Group) and Palmaz-Schatz® stents (Group)

	Group (n = 12)	Group (n = 12)	p
Reference diameter (mm)	2.97 ± 0.08	2.98 ± 0.09	0.668
Stented artery diameter (mm)	3.35 ± 0.18	3.34 ± 0.21	0.947
F/U MLD (mm)	2.50 ± 0.13	2.49 ± 0.19	0.897

1,000

Fig. 3

Core® stent PS® stent 10

통계 방법

±

Table 2. Histopathologic assessment of porcine coronary arteries stenting with Core® (Group) and Palmaz-Schatz® stent (Group)

	Group (n = 12)	Group (n = 12)	p
Injury score	1.58 ± 0.48	1.06 ± 0.82	0.096
Media area (mm²)	2.31 ± 1.14	2.18 ± 0.40	0.073
Intima area (mm²)	1.81 ± 0.67	2.93 ± 0.94	0.006
Intima/Media area ratio	0.91 ± 0.41	1.35 ± 0.40	0.023




Fig. 4. Pathologic findings showed smaller neointimal area in Core® stent (A, B) than in Palmaz-Schatz® stent (C, D).

Table 3. Measurement of angled changes after Core® (Group) and Palmaz-Schatz® stent(Group) im-plantation

	Group (n = 12)	Group (n = 12)	p
Pre-stent	19.6 ± 10.8	29.4 ± 21.2	0.47
Post-stent	8.8 ± 2.95	17.0 ± 9.67	0.22
Angled changes	13.2 ± 9.0	14.4 ± 11.1	0.88

Student 's t -
test p 0.05

결 과

12 PS® stent Core®
stent . PS® stent
6 , 5 , 1
Core® stent 6 , 4 ,
2
1) 1 2.
97 ± 0.08 mm, 2 2.98 ± 0.09 mm

1 3.35 ± 0.18 mm,
2 3.34 ± 0.21 mm
4 1 2.50 ± 0.13
mm, 2 2.49 ± 0.19 mm

(Table 1).
2) 1 1.58 ± 0.48,
2 1.06 ± 0.82 Core® stent Palmaz - Sch -
atz® stent

가 1 1.81 ±
0.67 mm², 2 2.93 ± 0.94 mm² Core® stent
Palmaz - Schatz® stent

(Table 2, Fig. 4).
3) Bending test Core® stent가 PS® stent
longitudinal straightening effect
가 (Fig. 5).

4) PS® stent Core® stent 가




Fig. 5. Comparisons of longitudinal straightening effect of Core® stent and PS® stent. Core® stent had more flexible property than PS® stent.

(Table 3).

고 찰

(elastic recoil)
1986 Sigwart
metallic stent
가 ,
8) ,
가 slotted tube
coil
가 coil
slotted tube radial force가
(acute stent recoil)
stent strut (tissue
prolapse) 9-11)
가 radial force
가
MAC® stent radial force
strut flexibility
vertical
tal strut radial force
horizon -

Core® stent strut MAC® strut , Core® stent가 PS® stent strai -
stent ghtening effect가
cell MAC® straighten -
stent flexibility trackability ing effect가 가
12)13)
strut radial
force Core® stent Core® stent가 PS® stent
longitudinal straightening
effect flexibility , bending test flexibility ,
PS® stent 가 가 Core® stent
가 , Core®
stent PS® stent 요 약
Core® 연구목적 :
stent PS® stent MAC® stent
Core® stent(E.O. Medical)
slotted tube stent Palmaz - Sc -
hatz® (PS) stent(Johnson & Johnson)
flexibility가
방 법 :
Core®
가 stent(1) PS® stent(2)
12 4
bending test
Core® stent conventional PTCA balloon
mount , 1 mm low crim -
ped profile longitudinal straightening effect
Core® stent PS® stent
4
stent
가 ,
Core® stent가 PS® stent
5
가 3 mm
minimal luminal diameter 가
2) Core® stent Palmaz -

Schatz® stent

1 1.81 ± 0.67 mm², 2 2.93 ± 0.94 mm² Core® stent Palmaz - Schatz® stent

3) Bending test Core® stent가 PS® stent

4) PS® stent Core® stent 가 (13.2 ±

9.0, 14.4 ± 11.1, p = 0.88).

결론 :

Core® stent가 PS® stent

, bending test flexibility , 가 Core® stent

가

중심 단어 :

REFERENCES

- 1) George BS, Voorhees WD 3rd, Roubin GS, Fearnot NE, Pinkerton CA, Raizner AE, et al. Multicenter investigation of coronary stenting to treat acute or threatened closure after percutaneous transluminal coronary angioplasty: Clinical and angiographic outcomes. *J Am Coll Cardiol* 1993;22:135-43.
- 2) Herrmann HC, Buchbinder M, Clemen MW, Fischman D, Goldberg S, Leon MB, et al. Emergent use of balloon-expandable coronary artery stenting for failed percutaneous transluminal coronary angioplasty. *Circulation* 1992;86:812-9.
- 3) Maiello L, Colombo A, Gianrossi R, McCanny R, Finzi

- L. Coronary stenting for treatment of acute or threatened closure following dissection after coronary balloon angioplasty. *Am Heart J* 1993;125:1570-5.
- 4) Fischman DL, Leon MB, Baim DS, Schatz RA, Savage MP, Penn I, et al. A randomized comparison of coronary stent placement and balloon angioplasty in the treatment of coronary artery disease. *N Eng J Med* 1994;331:496-501.
- 5) Serruys PW, de Jaegere P, Kiemeneij F, Macaya C, Rutsch W, Heyndrickx G, et al. A comparison of balloon-expandable stent implantation with balloon angioplasty in patients with coronary artery disease. *N Engl J Med* 1994;331:489-95.
- 6) Kim SH, Jeong MH, Jang YS, Bae Y, Kim JW, Cho JH, et al. Clinical experience of coronary MAC® stent. *Korean Circ J* 1998;28:1700-6.
- 7) Schneider JE, Berk BC, Gravanis MB, Santoian EC, Cipolla GD, Tarazona N, et al. Probucol decreases neointimal formation in a swine model of coronary artery balloon injury. A possible role for antioxidants in restenosis. *Circulation* 1993;88:628-37.
- 8) Sigwart U, Puel J, Mirkovitch V, Joffe F, Kappenberger L. Intravascular stents to prevent occlusion and restenosis after transluminal angioplasty. *N Engl J Med* 1987;316:701-6.
- 9) Lansky AJ, Roubin GS, O'Shaughnessy CD, Moore PB, Dean LS, Raizner AE, et al. Randomized comparison of GR-II® stent and Palmaz-Schatz® stent for elective treatment of coronary stenoses. *Circulation* 2000;102:1364-8.
- 10) Painter JA, Mintz GS, Wong SC, Popma JJ, Pichard AD, Kent KM, et al. Serial intravascular ultrasound studies fail to show evidence of chronic Palmaz-Schatz stent recoil. *Am J Cardiol* 1995;75:398-400.
- 11) Vrints CJ, Claeys MJ, Bosmans J, Conraads V, Snoeck JP. Effect of stenting on coronary flow velocity reserve: comparison of coil and tubular stents. *Heart* 1999;82:465-70.
- 12) Abhyankar DA, Luyue G, Bailey BP. Stent implantation in severely angulated lesions. *Cathet Cardiovasc Diagn* 1997;40:261-4.
- 13) Gyongyosi M, Yang P, Khorsand A, Glogar D. Longitudinal straightening effect of stents is an additional predictor for major adverse cardiac events. *J Am Coll Cardiol* 2000;35:1580-9.